

Number of double-normal pairs in space.

Andrey Kupavskii*

Abstract

Given a set V of points in $\mathbb{R}^d$, two points p, q from V form a double-normal pair, if the set V lies between two parallel hyperplanes that pass through p and q , respectively, and that are orthogonal to the segment pq . In this paper we study the maximum number $N_d(n)$ of double-normal pairs in a set of n points in $\mathbb{R}^d$. It is not difficult to get from the famous Erdős-Stone theorem that $N_d(n) = \frac{1}{2}(1 - 1/k)n^2 + o(n^2)$ for a suitable integer $k = k(d)$ and it was shown in the paper by J. Pach and K. Swanepoel that $\lceil d/2 \rceil \leq k(d) \leq d - 1$ and that asymptotically $k(d) \gtrsim d - O(\log d)$.

In this paper we sharpen the upper bound on $k(d)$, which, in particular, gives $k(4) = 2$ and $k(5) = 3$ in addition to the equality $k(3) = 2$ established by J. Pach and K. Swanepoel. Asymptotically we get $k(d) \leq d - \log_2 k(d) = d - (1 + o(1)) \log_2 k(d)$ and show that this problem is connected with the problem of determining the maximum number of points in $\mathbb{R}^d$ that form pairwise acute (or non-obtuse) angles.

1 Introduction

Let V be a set of n points in $\mathbb{R}^d$. We say that two points p, q from V form a *double-normal pair*, if the set V lies between two parallel hyperplanes H_p, H_q that pass through p and q , respectively, and that are orthogonal to the segment pq . We call a double-normal pair p, q *strict*, if $V \cap (H_p \cup H_q) = \{p, q\}$. Following this definition, we may consider the (strict) double-normal graph on V . We say that G is the *(strict) double-normal graph of V* if V is the

*Moscow Institute of Physics and Technology, École Polytechnique Fédérale de Lausanne; Email: kupavskii@yandex.ru Research supported in part by the Swiss National Science Foundation Grants 200021-137574 and 200020-14453 and by the grant N 15-01-03530 of the Russian Foundation for Basic Research.

vertex set of G and two points from V are connected by an edge iff they form a (strict) double-normal pair.

In 1957, Erdős [2] conjectured that if any pair of points in $V \subset \mathbb{R}^d$ forms a double-normal pair, then $|V| \leq 2^d$. This problem is better known in another, equivalent form: given a set $V \subset \mathbb{R}^d$ such that no three points from V form an obtuse angle, is it true that $|V| \leq 2^d$? This was proved by L. Danzer and B. Grünbaum in [1]. L. Danzer and B. Grünbaum in their paper asked the same question with a slight modification: all the triples in the set must form acute angles (instead of non-obtuse). In our terms, this question is the question of Erdős with double-normals changed to *strict* double-normals. L. Danzer and B. Grünbaum conjectured that the size of V , in which every two vertices form a strict double-normal pair, cannot be bigger than $2d - 1$. It was believed in for a long time, till Erdős and Füredi in [3] disproved it, showing an unexpectedly different bound, which states that the size of such V may be exponential in the dimension.

Formulated in double-normal graph terms, Erdős asked, how big a set $V \subset \mathbb{R}^d$ may be, if its double-normal graph is a complete graph. Or, what is the largest size $N(d)$ of a complete graph that may be represented as a double-normal graph in $\mathbb{R}^d$. Similarly, we can ask what is the largest size $N'(d)$ of a complete graph that may be represented as a *strict* double-normal graph in $\mathbb{R}^d$. In general, it is a classical question of extremal graph theory to ask what are the possible values of a clique number in a graph belonging to a certain class of graphs. In the paper [5] Martini and Soltan asked another classical extremal-graph-theory-type question for double normal graphs: what are the maximum numbers $N_d(n)$ and $N'_d(n)$ of double-normal pairs and strict double-normal pairs formed by pairs of points from a set among all sets of n points in $\mathbb{R}^d$? In terms of double-normal graphs this question is formulated as follows: what is the maximum number of edges in a (strict) double-normal graph on n vertices in $\mathbb{R}^d$?

It follows from the famous Erdős-Stone theorem that $N_d(n) = \frac{1}{2}(1 - 1/k)n^2 + o(n^2)$ for a suitable integer $k = k(d)$, where k is the largest integer such that for any $r \in \mathbb{N}$ we can find a double-normal graph in $\mathbb{R}^d$ that has a complete k -partite graph $K_k(r)$ with sizes of parts equal to r as a subgraph. Analogously, $N'_d(n) = \frac{1}{2}(1 - 1/k')n^2 + o(n^2)$ for a suitable integer $k' = k'(d)$. Note that $N_d(n) \geq N'_d(n)$ and $k \geq k'$. Thus, to find the asymptotic of $N_d(n)$ in case $N_d(n)$ being quadratic all we need is to determine k . The same takes place for $N'_d(n)$.

In the case of the plane and a two-dimensional sphere, however, these functions are linear in n . This was shown by J. Pach and K. Swanepoel in [8]. They, in fact, found exact values for the planar functions and their spherical analogues. In particular, they showed that $N'_2(n) = n$ and $N_2(n) = 3\lfloor n/2 \rfloor$.

In a complementary paper [7] they showed that already in $\mathbb{R}^3$ the functions are quadratic. More specifically, they showed that $k(3) = k'(3) = 2$. In general, they proved the following

Theorem 1 ([7]). *We have $\lceil d/2 \rceil \leq k'(d) \leq k(d) \leq d-1$ and asymptotically $k(d) \geq k'(d) \gtrsim d - O(\log d)$.*

Although Theorem 1 gives $k(3) = k'(3) = 2$, it does not allow to decide whether $k(4)$ or $k'(4)$ equals 2 or 3. In this paper we improve both the upper and the lower bounds from Theorem 1, which gives, in particular, $k(4) = k'(4) = 2$. While the improvement of the upper bound is easy to state, it is more difficult with the lower bound, since it is stated in Theorem 1 somewhat imprecisely. Thus, we first formulate a more accurate version of the asymptotical lower bound:

Theorem 2 ([7]). *Let $m \geq 2$. Suppose that there exist m points $p_1, \dots, p_m \in \mathbb{R}^d$ and m unit vectors $u_1, \dots, u_m \in \mathbb{R}^d$ such that, for all triples of distinct i, j, k , the angle $\angle p_i p_j p_k$ is acute, and*

$$\langle u_i, p_i - p_j \rangle < \langle u_i, p_k - p_j \rangle < \langle u_i, p_j - p_i \rangle.$$

Then, for any $N \in \mathbb{N}$, there exists a strict double-normal graph in $\mathbb{R}^{d+m}$ containing a complete m -partite $K_m(N)$. In particular, $k'(d+m) \geq m$.

Denote by $D''(m)$ the minimal dimension d in which points and vectors with conditions described in Theorem 2 exist. Then it is possible to state the result of Theorem 2 in the following form: $k'(d+m) + D''(k'(d+m)) \geq d+m$ or, changing the notation,

$$k'(d) + D''(k'(d)) \geq d. \quad (1)$$

We are almost ready to state both new lower and upper bounds. But before that we need to define two functions. Let $D(m)$ be the minimal dimension d in which m points forming only non-obtuse angles exist. Let $D'(m)$ be the analogous function for acute angles. We have $D(N(d)) = d, D'(N'(d)) = d$. It is also clear that $D(m) \leq D'(m) \leq D''(m)$. From the result of Danzer and Grünbaum we know that $D(m) = \lceil \log_2 m \rceil$, while from the result of V. Harangi [4], who improved the bound due to Erdős and Füredi to $N'(d) \geq 1.2^d$ for large d , we have $D'(m) \leq \log_{1.2} m$ for large m .

Theorem 3. *We have*

$$k(d) + D(k(d)) \leq d, \quad (2)$$

$$k'(d) + D'(k'(d)) \geq d. \quad (3)$$

Numerically, we have $k(d) + \log_2 k(d) \leq d$ for all d and $k'(d) + \log_{1.2} k'(d) \geq d$ for large d .

The upper bound from Theorem 3 together with the lower bound from Theorem 1 allows us to state the following corollary, which extends the result $k(3) = k'(3) = 2$ due to Pach and Swanepoel:

Corollary 1. *We have $k'(4) = k(4) = 2, k'(5) = k(5) = 3, k'(7) = k(7) = 4$.*

Thus, the next value to be determined is $k(6)$. In general, we ask if the bound (3) is sharp:

Problem 1. *Is it true that we have $k'(d) + D'(k'(d)) = d$ and, moreover, $k(d) + D'(k(d)) = d$?*

It would follow from a positive answer to Problem 1 that $k(d) = k'(d)$ for all d . We formulate this as a separate problem:

Problem 2. *Is it true that we have $k'(d) = k(d)$ for all d ?*

In Section 2 we present the proof of the upper bound from Theorem 3. In Section 3 we give the proof of the lower bound from Theorem 3.

We say that a graph G has a (strict) double-normal representation in $\mathbb{R}^d$, if there is a (strict) double-normal graph in $\mathbb{R}^d$ isomorphic to G . Instead of double-normal representations and strict double-normal representations one can consider *almost double-normal* representations. We say that a graph G admits an almost double-normal representation in $\mathbb{R}^d$, if for any $\delta > 0$ the graph may be represented in $\mathbb{R}^d$ by a set V of points, with two points x, y connected by an edge if for any $z \in V$ the angles $\angle xyz, \angle yxz \leq \frac{\pi}{2} + \delta$.

Denote by $k_1(d)$ the analogue of k, k' for this type of representations. In fact, slightly modifying the proof of both bounds in Theorem 3, one can prove

Theorem 4. *We have*

$$k_1(d) + D(k_1(d)) = d, \quad \text{that is, } k_1(d) + \lceil \log_2 k_1(d) \rceil = d.$$

Thus, we can determine the exact value of the k -function in this somewhat modified scenario. We leave it to the reader to check that the proof of theorem 3 works for this situation.

Throughout the paper we use the following notations. The scalar product of two vectors u, v is denoted by $\langle u, v \rangle$. The linear span of a set of vectors U we denote $\text{lin } U$, the affine span of a set of points S we denote $\text{aff } S$. By $\dim W$ we denote the dimension of a subspace or a plane W . The angle formed by points x, y, z with sides yx, yz and its angular measure we denote by $\angle xyz$. Throughout the paper all angles have angular measure from $(0, \pi)$. The Euclidean distance between two points x, y we denote $\|x - y\|$.

2 Upper bound on $k(d)$

We follow the steps of the proof from [7] used to obtain the upper bound on $k(d)$. The goal is to prove a strengthened version of Proposition 8 from [7], which is formulated in the forthcoming theorem. Despite that some parts of the proof of the theorem are identical to the ones from [7], we will present an almost complete proof of the theorem for the sake of clarity of exposition. In what follows we do not specify the parts that are taken from [7], but after the proof we are going to comment on what is new and what is old in the statement and in the proof.

Theorem 5. *Put $k = k(d)$. There exists a family of (not necessarily distinct) vectors u_{ij} , where $i, j = 1, \dots, k$ so that the following conditions hold:*

$$U_1 := \{u_{11}, \dots, u_{kk}\} \text{ form an orthogonal set of vectors.} \quad (4)$$

$$u_{ii} \text{ is orthogonal to } u_{jl} \text{ for any } i, j, l = 1, \dots, k, \text{ such that } j \neq l. \quad (5)$$

$$\dim \operatorname{lin} U_2 \geq \log_2 k \text{ for } U_2 := \{u_{sl} : s, l = 1 \dots, k, s \neq l\}. \quad (6)$$

Remark. Bound (2) from Theorem 3 follows easily from Theorem 5 by the following observation: $d \geq \dim \operatorname{lin}(U_1 \cup U_2) = \dim \operatorname{lin} U_1 + \dim \operatorname{lin} U_2 \geq k + \lceil \log_2 k \rceil = k + D(k)$.

The rest of the section is dedicated to the proof of Theorem 5. In what follows, we will use the following notation: $f(\epsilon) \gg g(\epsilon)$, if $\lim_{\epsilon \rightarrow 0} g(\epsilon)/f(\epsilon) \rightarrow 0$, in addition to the traditional (o, O, Ω) -notation with respect to the limit $\epsilon \rightarrow 0$. Also, we will say that two vectors are ϵ -orthogonal (ϵ -parallel), if the angle between them is equal to $\pi/2 + f(\epsilon)$ ($f(\epsilon)$), where $f(\epsilon) = o(1)$.

First we give a sketch of the proof.

Step 1. We show that for any $\epsilon > 0$ there exists N such that for any complete k -partite graph $K_k(N)$ with parts $V_1, \dots, V_k$ of size N contained in a double-normal graph and for any i we can choose points A_i, a_i, b_i, C_i out of V_i such that the following conditions hold:

$$\text{Points from } \{A_i, a_i, b_i, C_i\} \text{ satisfy } \angle A_i a_i b_i \geq \pi - \epsilon, \angle a_i b_i C_i \geq \pi - \epsilon, \quad (7)$$

$$\|A_{i+1} - C_{i+1}\| \leq \epsilon \|A_i - C_i\|, \quad (8)$$

$$\|A_i - b_i\| \geq 1/2 \|A_i - C_i\|, \quad (9)$$

$$\|a_i - b_i\| \leq \epsilon \|A_i - b_i\|. \quad (10)$$

Step 2. For the points chosen in Step 1 and sufficiently small $\epsilon > 0$, set $u_{ii} = \|A_i - b_i\|^{-1}(A_i - b_i)$ and $u_{ij} = \|b_j - b_i\|^{-1}(b_j - b_i)$. We verify the

properties

$$|\langle u_{ii}, u_{ij} \rangle| \leq \epsilon, i, j = 1, \dots, k, i \neq j, \quad (11)$$

$$|\langle u_{ii}, u_{jj} \rangle| \leq 4\epsilon, \quad i, j = 1, \dots, k, i \neq j, \quad (12)$$

$$|\langle u_{ii}, u_{jl} \rangle| = o(1), \quad i, j, l = 1, \dots, k, i \neq j \neq l, i \neq l. \quad (13)$$

Step 3. We set $\epsilon = 1/n$ and take a subsequence of n 's such that each $u_{ij}^{(n)}$ converge as $n \rightarrow \infty$. The limiting set of vectors is the desired one.

Step 4. We give the estimate on the dimension of $\text{lin } U_2$.

Remark. Note that we have the same notation for the vectors for a fixed ϵ and the limiting vectors, which, however, should not cause confusion since it should be clear from the context which out of the two we are dealing with. In Step 1, Step 2 we deal only with u_{ij} for a fixed ϵ .

Step 1. First, choose some $0 < \epsilon < 1$ and take a sufficiently large N depending on ϵ and d and apply [Theorem 2.6, [6]] to each part V_i of the complete multipartite graph $K_k(N)$ contained in a double-normal graph. We get that any V_i contains a subset V'_i of points $\{p_i^1, \dots, p_i^{N'}\}$ such that for any $i < j < l, i, j, l = 1, \dots, N'$ we have $\angle p_i p_j p_l \geq \pi - \epsilon$, where $N' = (2t + 3)t^{k-1}$ and $t = \lceil (\epsilon \cos \epsilon)^{-1} \rceil$. We replace the original V_i by V'_i .

Next, we modify the sets V_i so that condition (8) holds. We do it in k steps. At the first step we rename the parts in such a way that V_1 has the largest diameter over all V_i . At step j we do the following to each $V_i, i = j + 1, \dots, k$. Take $V_i, i > j$ and project all the points of V_i onto the segment that connects the furthest points of V_i and split it into t parts of equal length. By the pigeon-hole principle, there is a part that has at least $\frac{1}{t}$ -th fraction of all the points in V_i . Denote it by V'_i . The diameter of V'_i is at most $\frac{1}{t \cos \epsilon}$ multiplied by the diameter V_i , where $\frac{1}{t \cos \epsilon} \leq \epsilon$. The inequality on the diameter V'_i holds since any segment connecting points of V_i form an angle at most ϵ with the diameter. After that we change V_i for V'_i and rename them in such a way that V_{j+1} has the largest diameter over $V_i, i > j$.

Note that at the end of this procedure V_i has at least $N't^{-i+1} \geq 2t + 3$ points. As A_i and C_i we choose the points that form a diameter of V_i . Thus, the condition (8) is satisfied.

To obtain (10) for some vertices in V_i , we again project the points of V_i on $A_i C_i$ and split the segment into parts of length $\frac{1}{2t} \|A_i - C_i\|$. By the pigeon-hole principle, there will be a part that contains at least two (projected) points apart from A_i and C_i . Denote them by b_i and a_i . We have $\|b_i - a_i\| \leq \frac{1}{2t} \|A_i - C_i\|$ and either $\min\{\|A_i - b_i\|, \|A_i - a_i\|\} \geq \frac{1}{2} \|A_i - C_i\|$ or $\min\{\|C_i - b_i\|, \|C_i - a_i\|\} \geq \frac{1}{2} \|A_i - C_i\|$. Rename the points in such a way that

they appear on $A_i C_i$ in the order A_i, a_i, b_i, C_i and $\|A_i - b_i\| \geq \frac{1}{2}\|A_i - C_i\|$. Now consider the preimages of a_i, b_i before the projection. We also denote them by a_i, b_i . We have $\|a_i - b_i\| \leq \frac{1}{t \cos \epsilon} \|A_i - b_i\| \leq \epsilon \|A_i - b_i\|$. Therefore, (10) is verified for A_i, a_i, b_i, C_i . Moreover, (9) is verified as well. This completes Step 1.

Step 2.

The proof of (11) and (12) is identical to the one presented in [7] and can be losslessly separated from the rest of the proof. Thus, we omit it, only mentioning that it uses properties (7), (8), (9) of the configuration.

The rest of Step 2 is devoted to verifying (13). We need to prove that for any pairwise distinct $i, j, l = 1, \dots, k$ the vectors $A_i - b_i$ and $b_l - b_j$ are ϵ -orthogonal. For simplicity put $i = 1, j = 2, l = 3$. Consider the two two-dimensional planes $\gamma_a = \text{aff}\{b_2, b_1, a_1\}$ and $\gamma_b = \text{aff}\{b_3, b_1, a_1\}$.

Observation 1. *Consider the projections b_2^γ, b_3^γ of points b_2, b_3 on the planes γ_b, γ_a respectively. Then either b_2^γ lies to the same side from the line $b_1 b_3$ as a_1 or b_3^γ lies to the same side from the line $b_1 b_2$ as a_1 .*

Proof. Consider projections b'_2, b'_3 of b_2, b_3 on the line $a_1 b_1$. Since $\angle a_1 b_1 b_i \leq \pi/2$, $i = 2, 3$, both b'_2, b'_3 lie in the same closed halfline emanating from b_1 as a_1 . Suppose that b'_3 is at least as far away from b_1 as b'_2 . We then claim that b_3^γ lies to the same side from the line $b_1 b_2$ as a_1 .

The point b_3^γ lies on a line l that is orthogonal to $a_1 b_1$ and that passes through b'_3 . We also know that b_3^γ must lie between the two lines that are orthogonal to $b_2 b_1$ and that pass through the points b_2 and b_1 . Otherwise, the pair b_1, b_2 is not double-normal. Then it is easy to see that the part of l that is between the two lines does not cross $b_2 b_1$ and thus lies in the same closed halfplane as a_1 . $\square$

Consider a unit vector v_a that is lying in the plane γ_a , orthogonal to $b_2 b_1$ and pointing to the side of $b_1 b_2$ on which a_1 lies. We remark that v_a is ϵ -parallel to u_{11} . In what follows, we w.l.o.g. assume that b_3^γ lies to the same side from the line $b_1 b_2$ as a_1 . Or, in other words, that $\langle u_{23}, v_a \rangle \geq 0$. Consider the projection b_3^γ of b_3 on the plane γ_a . We are going to analyze possible locations of b_3^γ using the fact that almost all pairs of points from $\{b_1, b_2, b_3, a_1, A_1\}$ form a double normal (except for the pairs formed by $\{a_1, A_1, b_1\}$). We denote the projection of A_1 on γ_a by A_1^γ . See Figure 1, that depicts the plane γ_a . Note that all the points on the figure indeed lie in that plane, while, in general, A_1 does not lie in γ_a .




Figure 1

First, as we have already mentioned, since b_1b_2 is a double-normal pair, b_3^γ lies in a strip bounded by the two lines that are orthogonal to b_1b_2 and that run through b_1 and b_2 respectively (see Figure 2). Second, b_3^γ lies in a strip bounded by b_1b_2 itself and a line that is parallel to b_1b_2 and that passes through a_1 . One part of this condition holds since $\langle u_{23}, v_a \rangle \geq 0$, while the other one holds since otherwise one of the angles $\angle b_3a_1b_2$ or $\angle b_3a_1b_1$ would be greater than $\pi/2$, which, in turn, gives that b_3a_1 is not a double-normal pair. Thus, b_3^γ falls into the hatched rectangle.

For simplicity, we rescale the configuration so that $\|b_2 - b_1\| = 1$. Next, we consider two cases.

Case 1 The segment A_1b_1 is “not much bigger” than b_2b_1 : $\|A_1 - b_1\| = O(\epsilon^{-1/3})$.



If b_3 projects into the quadrangle, then

The inequality above holds since b_3^γ falls in the angular domain $Q_1 b_2 b_1$, so $\angle b_3^\gamma b_2 b_1 \leq \angle Q_1 b_2 b_1$. The pre-last equality holds since $\|Q_2 - b_1\| \leq \|a_1 - b_1\| \leq \epsilon \|A_1 - b_1\| = O(\epsilon^{2/3})$. Thus, the vector $b_3^\gamma - b_2$ is ϵ -parallel to $b_1 - b_2$, which means that $b_3^\gamma - b_2$ and, consequently, u_{23} is ϵ -orthogonal to v_a and to u_{11} .

Lemma 1. *Consider a circle S on the plane with the center in O and with a point B on the circle (See Figure 3). Take a point A inside the circle, such that AB is ϵ -parallel to OB and $\|A - B\| = o(\|O - B\|)$. Draw a line l through A that is orthogonal to AB . Then we have $\|A - B\| = o(\min\{\|P_1 - A\|, \|P_2 - A\|\})$, where P_1, P_2 are the intersection points of l and S .*

Proof. Since the triangle ABP_1 is similar to AP_2B' , we have

$$\frac{\|A - B\|}{\|A - P_1\|} = \frac{\|A - P_2\|}{\|A - B'\|} = \frac{\|A - P_2\|}{2\|O - B\| \cos \angle OBA - \|A - B\|} < \frac{\|P_1 - P_2\|}{2\|O - B\| \cos \angle OBA - \|A - B\|} \rightarrow 0 \quad \text{as} \quad \frac{\|A - B\|}{\|O - B\|} \rightarrow 0.$$

Similar reasoning applies to $\|A - P_2\|$. □




Figure 3

The following fact we use several times, so we formulate it as an observation:

Observation 2. *If $\|b_3 - b_3^\gamma\| \gg \|b_3^\gamma - b_2\|$, then u_{23} is ϵ -orthogonal to u_{11} .*

In the remaining triangular cases, according to Observation 2, it is enough to verify the assumption of the observation. Assume that b_3^γ falls in the triangle Yb_2Q_1 and it lies on the line r that passes through b_2 . In the space $\mathbb{R}^d$ take a two-dimensional plane π_r that contains r and b_3 . It is orthogonal to the plane γ_a . Denote by S the (full-dimensional) sphere in $\mathbb{R}^d$ that has A_1b_2 as a diameter. Finally, consider a circle $S_r = \pi_r \cap S$ in the plane π_r . Denote its center by O and the projection of O on the plane γ_a by O^γ .

By the choice of A_1, a_1, b_1 we have $\|O - O^\gamma\| \leq \|A_1 - A_1^\gamma\| = O(\epsilon \|A_1 - b_1\|)$. Denote by P_1 and P_2 (from Figure 2) are the projections of A_1 (or A_1^γ) on

Yb_2 and b_1b_2 respectively. At the same time,

$$\|O - b_2\| \geq \frac{1}{2} \begin{cases} \|P_1 - b_2\| & \text{if } b_3^\gamma \in XYb_2, \\ \|P_2 - b_2\| & \text{if } b_3^\gamma \in XQ_1b_2 \end{cases} = \Omega(\min\{\|A_1 - b_1\|, 1\}). \quad (14)$$

The inequality above holds due to the following. All the points x of the segments $A_1^\gamma P_1, A_1^\gamma P_2$ lie inside the sphere S since the angle A_1xb_2 is not acute. Thus, the intersection of r and these segments lie inside the circle S_r and the distance from the intersection point to b_2 is less than the diameter of S_r . On the other hand, any point on $A_1^\gamma P_i$ is farther away from b_2 than P_i . The equality in (14) holds since $\|P_2 - b_1\| = O(\epsilon\|A_1 - b_1\|) = O(\epsilon^{2/3})$.

Therefore,

$$\frac{\|O - O^\gamma\|}{\|O - b_2\|} \leq \frac{O(\epsilon\|A_1 - b_1\|)}{\Omega(\min\{\|A_1 - b_1\|, 1\})} = O(\epsilon^{2/3}).$$

It means that Ob_2 and $O^\gamma b_2$ are ϵ -parallel, and, consequently, Ob_2 and $b_3^\gamma b_2$ are ϵ -parallel.

Since b_3 and b_2 form a double-normal pair, $\angle b_2b_3A_1 \leq \pi/2$, which means that point b_3 lies on the outside of the (open) ball bounded by the sphere S , or, in other words, in the plane π_r the point b_3 on the outside of the circle S_r . We also note that both P_1 and P_2 lie on S .

Assume for a second that we proved that $\|b_3^\gamma - b_2\| = o(\|O - b_2\|)$. Then we apply Lemma 1 and obtain that $\|b_3 - b_3^\gamma\| \gg \|b_3^\gamma - b_2\|$, which is the desired statement. So it is enough to show for both triangles XYb_2 and Xb_2Q_1 that the radius of S_r is “much bigger” than $\|b_3^\gamma - b_2\|$.

Consider first XYb_2 . Suppose that r intersects XY and $A_1^\gamma P_1$ in the points R_1 and R_2 respectively. Both R_i are inside S_r , so the leftmost point of intersection of S_r and r lies to the left from $A_1^\gamma P_1$ (or coincides with one of P_1, A_1^γ), thus, $2\|O - b_2\| \geq \|R_2 - b_2\|$. Therefore,

$$\|b_3^\gamma - b_2\| \leq \|R_1 - b_2\| = O(\epsilon\|R_2 - b_2\|) = o(\|R_2 - b_2\|) = o(\|O - b_2\|),$$

where the first equality holds because of the choice of A_1, a_1, b_1 .

Consider the case of the triangle Xb_2Q_1 . On the one hand, in this case we have

$$\|b_3^\gamma - b_2\| \leq \|Y - Q_1\| + \|Y - b_2\| = O(\epsilon^{1/2}).$$

On the other hand, by (14) we have $\|O - b_2\| = \Omega(1)$. This again gives $\|b_3^\gamma - b_2\| = o(\|O - b_2\|)$.

Case 2 The segment A_1b_1 is “much bigger” than b_2b_1 : $\|A_1 - b_1\| \gg \epsilon^{-1/3}$.




Figure 4

This situation is depicted on Figure 4. The point P from Figure 4 is chosen in the following way. First, $\angle Qb_2b_1 = \epsilon^{1/6}$, and, second, $\angle A_1^\gamma Pb_2 = \pi/2$. The last condition is equivalent to saying that P lies on the sphere with diameter A_1b_2 . The point Q is the intersection of the line Pb_2 and the segment Q_2b_1 or the segment Q_2X .

In this case we again have three differently hatched regions, in which the point b_3 may be projected. On the figure these are two triangles YXb_2 and Qb_2b_1 , and a quadrangle Xb_2Q_2 . However, the line b_2P may not intersect the segment Q_2b_1 and intersect Q_2X instead.

The argument for the situation when b_3 projects inside XYb_2 is identical to the one for that situation in Case 1.

The vectors $P - b_2$ and $b_1 - b_2$ are ϵ -parallel, so if the point b_3^γ lies in the angular domain bounded by b_2Q and b_2b_1 (the triangle Qb_2b_1 in Figure 4), then $b_3^\gamma - b_2$ is ϵ -parallel to $b_1 - b_2$ and, thus, u_{32} and u_{11} are ϵ -orthogonal.

The last case to consider is when b_3^γ falls inside the angular domain bounded by $A_1^\gamma b_2$ and Pb_2 (the horizontally hatched quadrangle on Figure 4). Arguing as in Case 1, consider the circle S_r . We again show that the radius $\|O - b_2\|$ of S_r is “much bigger” than $\|b_3^\gamma - b_2\|$. On the one hand, we have

$$\|b_3^\gamma - b_2\| \leq \|b_1 - b_2\| + \|Y - b_2\| = O(1 + \epsilon\|A_1 - b_1\|).$$

On the other hand, using the same argument as the one used for the inequality in (14) we see that the diameter of S_r is at least $\|P - b_2\|$, and

$$\|P - b_2\| \geq \sin \angle Pb_2b_1 \|A_1^\gamma - b_2\| = \Omega(\epsilon^{1/6}\|A_1 - b_2\|) = \Omega(\epsilon^{1/6}\|A_1 - b_1\|).$$

Thus, $\|O - b_2\| = \Omega(\epsilon^{1/6}\|A_1 - b_1\|)$. Combined with the condition that $\|A_1 - b_1\| \gg \epsilon^{-1/3}$ we get that the $\|b_3^\gamma - b_2\| = o(\|O - b_2\|)$. At the same time, as in Case 1 we have $\|O - O^\gamma\| = O(\epsilon\|A_1 - b_1\|)$ and therefore

$$\frac{\|O - O^\gamma\|}{\|O - b_2\|} = O(\epsilon^{5/6}).$$

Therefore, Ob_2 and its projection on the plane γ_a are ϵ -parallel. Thus, we can apply Lemma 1 and get that $\|b_3 - b_3^\gamma\| \gg \|b_3^\gamma - b_2\|$. By Observation 2 this concludes the proof.

Step 3. We set $\epsilon = 1/n$ and choose $A_i^{(n)}, a_i^{(n)}, b_i^{(n)}, c_i^{(n)}$ as in the first part of the proof. All vectors $u_{ij}^{(n)}$ lie on a unit sphere, which is a compact space. Thus, passing to a subsequence, we can assume that for each i, j the sequence $u_{ij}^{(n)}$ converges to, say, u_{ij} . It is easy to see from the properties (11), (12), (13) that $\langle u_{ii}, u_{jj} \rangle = 0$ and that $\langle u_{ii}, u_{jl} \rangle = 0$ for any $i, j, l = 1, \dots, k$ if $j \neq l$. Thus, we have verified parts (4) and (5) of Theorem 5.

Step 4. This is the last part of the proof, in which we are going to obtain the bound on the dimension of $\text{lin } U_2$. To bound $\dim \text{lin } U_2$, we want to work with points instead of vectors. All u_{ij} with $i \neq j$ come as a limit of scaled $b_j^{(n)} - b_i^{(n)}$. We choose some $\delta > 0$ and n sufficiently large so that the angle between $u_{ij}^{(n)}$ and u_{ij} is less than $\delta/2$ for any $i \neq j$. Next, we project all the points $b_i^{(n)}$ on the subspace $\text{lin } U_2$, which is interpreted as an affine plane passing through zero. Denote by $B_i^{(n)}$ the corresponding projections. They are obviously distinct for different points if δ is small enough. Otherwise, if $B_i^{(n)} = B_j^{(n)}$ for some $i \neq j$, then $u_{ij}^{(n)}$ is orthogonal to U and, consequently, to u_{ij} .

The angle between $B_j^{(n)} - B_i^{(n)}$ and $u_{ij}^{(n)}$ is less than the angle between $u_{ij}^{(n)}$ and u_{ij} (since $B_j^{(n)} - B_i^{(n)}$ is a scaled projection of $u_{ij}^{(n)}$ on U). Therefore, by the triangle inequality, for any pairwise distinct $i, j, l = 1, \dots, k$ we have $\angle B_i^{(n)} B_j^{(n)} B_l^{(n)} \leq \angle b_i^{(n)} b_j^{(n)} b_l^{(n)} + \alpha_1 + \alpha_2 \leq \pi/2 + \alpha_1 + \alpha_2$, where α_1 is the angle between vectors $u_{ji}^{(n)}$ and $B_i^{(n)} - B_j^{(n)}$ and α_2 is the angle between vectors $u_{jl}^{(n)}$ and $B_l^{(n)} - B_j^{(n)}$. Both α_1 and α_2 are less than $\delta/2$, so for any pairwise distinct $i, j, l = 1, \dots, k$ we have $\angle B_i^{(n)} B_j^{(n)} B_l^{(n)} \leq \pi/2 + \delta$.

The above considerations allowed us to construct an affine plane of dimension $\dim \text{lin } U_2$ and a set of k points on this plane without triples that form an angle bigger than $\pi/2 + \delta$ for some $\delta > 0$. Moreover, we may make δ as small a positive number as we want it to be. It turns out that $k \leq 2^{\dim \text{lin } U_2}$, which is stated in

Lemma 2. *For any natural d there exists a $\delta > 0$ such that the following holds. Consider any k points $p_1, \dots, p_k$ in $\mathbb{R}^d$ such that for any distinct $i, j, l \in \{1, \dots, k\}$ we have $\angle p_i p_j p_k \leq \pi/2 + \delta$. Then $k \leq 2^d$.*

Remark. The statement of Lemma 2 for $\delta = 0$ is a result by L. Danzer and B. Grünbaum [1]. A statement equivalent to Lemma 2 is mentioned without a proof in [3], so it may be well known. However, in [3] it looks like the authors somewhat inaccurately attributed Lemma 2 to Danzer and Grünbaum. Thus, we present the proof of the lemma here for completeness. We remark that, contrary to intuition, it does not follow from the result due to Danzer and Grünbaum via a direct compactness argument.

Proof. We prove the statement by induction. It is easy to check for the plane. Now suppose we have proved it for the dimension $d - 1$ and pass on to the dimension d . Assume that the lemma is not true in $\mathbb{R}^d$. Then there exists a sequence of point sets $W_j = \{p_1^j, \dots, p_k^j\}$ in $\mathbb{R}^d$ so that $\|W_j\| = k = 2^d + 1$ and the maximal angle in W_j is less than $\pi/2 + 1/j$. Scale and translate these sets so that all W_j have unit diameter and so that all of them lie in a unit ball. Passing to subsequences, we have convergent sequence, which gives a limiting multiset of points $W = \{p_1, \dots, p_k\}$. Naturally, in W all the angles are non-obtuse and W is of unit diameter. If none of p_i coincide, then we get a contradiction with the result by Danzer and Grünbaum. Thus, at least two points among p_i must coincide.

Assume that the maximum multiplicity of an element in W is m , while the corresponding point in W is, w.l.o.g., p_1 . Denote the subset of points that participate in the sequences that do converge to p_1 in every W_j by M_j . Since W is of unit diameter, all of the points in W cannot coincide and, thus, $m \leq k - 1 = 2^d$. Choose an integer l so that $2^{l-1} < m \leq 2^l$, $1 \leq l \leq d$. Consider the set of vectors U_j : $U_j = \{u_{sr}^j = p_r^j - p_s^j : p_r^j, p_s^j \in M_j, r \neq s\}$ and a limiting set of vectors U . We claim that, first, $\dim \text{lin } U \geq l$ and, second, $\text{aff } W$ is orthogonal to $\text{lin } U$ in the sense that any vector formed by points from $\text{aff } W$ is orthogonal to any vector from $\text{lin } U$.

To show that $\dim \text{lin } U \geq l$ we have to apply induction. We choose sufficiently large j so that the vectors from U_j are close to the limiting vectors and, as in Step 4, project all points from M_j on $\text{lin } U$. We get angles as close to $\pi/2$ as we want to (it depends on j only), so for a right choice of j we can apply induction to M_j and $\text{lin } U$ and get the stated inequality.

To show that W spans a plane that is orthogonal to $\text{lin } U$, we assume the contrary. Choose a point from W , say, p_2 , such that the vector $p_2 - p_1$ is not orthogonal to $\text{lin } U$. This means that there is a pair of indices s, r

and $\epsilon > 0$, so that $p_2 - p_1$ and u_{sr} form an angle at least $\pi/2 + 4\epsilon$ big. If we choose a sufficiently large j , then $p_2 - p_s$ and $p_2^j - p_s^j$ form an angle α_1 at most ϵ , similarly, $p_r - p_s$ and $p_r^j - p_s^j$ form an angle α_2 at most ϵ and, finally, there is no angle bigger than $\pi/2 + \epsilon$ in W_j . But then we arrive at a contradiction, since on the one hand, $\angle p_r^j p_s^j p_2^j \leq \pi/2 + \epsilon$, on the other hand, $\angle p_r^j p_s^j p_2^j \geq \angle p_r p_s p_2 - \alpha_1 - \alpha_2 \geq \pi/2 + 2\epsilon$.

Therefore, W has at least $\frac{2^d}{m} + 1 \geq 2^{d-l} + 1$ distinct points and $\dim \text{aff } W \leq d - \dim \text{lin } U \leq d - l$. We may apply induction and arrive to a contradiction. $\square$

This proves (6) and completes the proof of Theorem 5.

Remark. As we have promised, we discuss here the differences between Theorem 5 and Proposition 8 from [7].

In the formulation (and in essence) the main difference is that in Theorem 5 we have a much more general condition (5) rather than the condition “ u_{ii} and u_{ij} are orthogonal, if $i \neq j$ ” from Proposition 8. We also have the condition (6) that bounds the dimension of $\text{lin } U_2$. This, as well as Step 4, is not present in the proposition.

Step 1 in our proof is really similar to Step 1 from the proposition. A large part is just identical to the analogous part from the proposition. However, we choose 4 points (with specific distances) instead of 3 in the proposition, so the last paragraph of Step 1 is new.

The parts of Step 2 that are presented (recall that we omit the proof of (11) and (12)) are completely new.

Step 3 is really similar to that step in the proposition, however, it is shorter, since we are working only with vectors.

As we have already mentioned, there is no analogue of Step 4 in the paper [7].

3 Lower bound on $k'(d)$

The proof of (3) is based on a slight modification of the proof of Theorem 2. In this section we first give a complete proof of (3) and then the sketch of the original proof of Pach and Swanepoel. In this way the reader would have the chance to compare the two proofs. We split the section into two subsections.

3.1 Proof of (3)

Step 1. Fix $p_1, \dots, p_m \in \mathbb{R}^d$ so that they form only acute angles. Thus, they are in convex position. For each p_i choose a hyperplane H_i that touches

the convex hull of $\{p_1, \dots, p_m\}$ only in p_i and a normal vector u_i to H_i , that points towards the convex hull. Embed $\mathbb{R}^d$ as a plane Γ in $\mathbb{R}^{d+m}$. For each p_i consider a two-dimensional plane Π_i spanned by u_i and v_i , where v_i are orthogonal to Γ and are mutually orthogonal as well.

Step 2. For every i do the following. Project all p_j on Π_i (they all fall on a half-line that starts at p_i and goes in the direction of u_i). Choose point c_i on the half-line so that the circle C_i in Π_i with center in c_i and that passes through p_i contains strictly inside all the projections of p_j . Denote the other intersection of the halfline and the circle by q_i (See Fig. 5, where we omitted the subscript indices i for simplicity). Choose an ϵ and a segment I_i of the halfline in such a way that I_i is disjoint with the ϵ -neighborhoods of the points p_i, q_i , but contains each p'_j with an ϵ -neighborhood (on the halfline). The endpoints of I_i are denoted by a and b .




Figure 5

Step 3. In this step we construct the vertices of the (supergraph of a) complete multipartite graph $K_m(N)$. Choose points $x_1^i, \dots, x_N^i \in C_i$ in the following way. First, for any z that belongs to the segment I_i and for any $s, l \in \{1, \dots, N\}, s \neq l$, we have $\angle z x_s x_l < \pi/2$. Second, all x_s are in ϵ -neighborhood of p_i . Put $V_i = \{x_1^i, \dots, x_N^i\}$.

In what follows we justify that such a choice is possible. For simplicity, for the rest of Step 3 we drop all the subscripts (and superscripts) i that refer to the part V_i . The second condition is very easy to satisfy, thus, we only have to show how to choose x_s such that for any $z \in I$ and for any different $s, l = 1, \dots, N$ we have $\angle zx_s x_l < \pi/2$. Recall that the endpoints of I are denoted by a, b (see Figure 5). We describe the procedure of choosing x_s recursively. We take as x_1 an arbitrary point from the upper semicircle that is ϵ -close to p (but, obviously, different from p). Next, having chosen $x_1, \dots, x_s$, we choose x_{s+1} from the arc px_s so that all the angular constraints are satisfied. It is easy to see that for any $0 \leq k \leq s-1$ and any point z between a and b we have $\angle x_{s+1}x_{s-k}z \leq \angle x_{s+1}x_s z$ and $\angle x_{s-k}x_{s+1}z \leq \angle x_s, x_{s+1}, z$. Moreover, $\angle x_s x_{s+1} z \leq \angle x_s x_{s+1} a$ and $\angle x_{s+1} x_s z \leq \angle x_{s+1} x_s b$. Therefore, we need to satisfy only two constraints: $\angle x_{s+1} x_s b$ is acute, and $\angle x_s x_{s+1} a$ is acute.

To satisfy the first constraint, we draw a chord from x_s that is orthogonal to bx_s and choose x_{s+1} from an open arc connecting p and the second intersection point of the chord.

Next we explain how to satisfy the second constraint. Assume we made a choice of x_{s+1} . Consider a point $x'_s \in C$, which is symmetric to x_s with respect to the center c . Next, consider a chord that contains x_{s+1} and a . Its other intersection with C denote by y_{s+1} . If y_{s+1} lies on an arc $x'_s q$, then $\angle x_s x_{s+1} a$ is acute. Thus, it suffices to choose x_{s+1} on the open arc $a_s p$, where a_s is the second intersection point of the chord that passes through x'_s and a .

It is clear that we can satisfy both restrictions if we choose x_{s+1} close enough to p .

Step 4. In this step we verify that the strict double-normal graph on the set of vertices $\cup_i V_i$ contains $K_m(N)$ as a subgraph. We need to show that any two points y, z from different V_i form a strict double normal. For that we need to do a case analysis. We assume throughout that ϵ is chosen to be sufficiently small. Take $x, y, z \in \cup_i V_i$. If x, y, z are all from different parts V_i, V_j, V_k , then $\angle xyz$ is acute since points x, y, z are ϵ -close to p_i, p_j, p_k and $\angle p_i p_j p_k$ is acute. If $x, z \in V_i, y \in V_j$, then $\|x - z\|$ is small compared to $\|x - y\|, \|z - y\|$, so the angle $\angle xyz$ is small and, thus, acute. The last case is if $x, y \in V_i$ and $z \in V_j$. Then $\angle xyz$ is acute since the angle $\angle xyz'$ is acute, where z' is a projection of z on Π_i . In turn, since z' is in the ϵ -neighborhood of p'_j , we have $z' \in I_i$. Therefore, Step 3 tells us that $\angle xyz'$ is acute. Thus, $\cup_i V_i$ indeed contains $K_m(N)$ as a subgraph.

3.2 Sketch of the proof of Theorem 2

Step 1. Fix $p_1, \dots, p_m \in \mathbb{R}^d$ and $u_1, \dots, u_m \in \mathbb{R}^d$ as in the assumption of the theorem. Embed $\mathbb{R}^d$ as a plane Γ in $\mathbb{R}^{d+m}$. For each p_i consider the two-dimensional plane Π_i spanned by u_i and v_i , where v_i are orthogonal to Γ and are mutually orthogonal as well.

Step 2. For every i do the following. Project all p_j on Π_i (they all fall on a half-line that starts at p_i and goes in the direction of u_i). Choose point c_i on the half-line so that the circle C_i in Π_i with center in c_i and that passes through p_i has all the projections of p_j inside an open segment $c_i q_i$, where q_i is the intersection of the halfline and the circle, which is different from p_i . By I_i denote the segment $c_i q_i$, out of which we exclude ϵ -neighborhoods of c_i, q_i for some small $\epsilon > 0$. Make sure that ϵ is so small that p'_i with their ϵ -neighborhood fall inside I_i .

Step 3. Choose points $x_1^i, \dots, x_N^i \in C_i$ in the following way. First, for any z that belongs to the segment I_i and for any different $j, l = 1, \dots, N$ we have $\angle z x_j x_l < \pi/2$. Second, all x_i are in ϵ -neighborhood of p_i . Put $V_i = \{x_1^i, \dots, x_N^i\}$.

Step 4. This step repeats word for word Step 4 from the previous section (or, rather, the way around: Step 4 in the proof of (3) is the same as Step 4 in this proof).

Remark. The modifications that lead to the proof of the lower bound in Theorem 3, actually simplify significantly the proof of the lower bound as stated in Theorem 1. It is due to the fact that we may use an Erdős-Füredi-type bound right away, while in [7] the authors had to consider and analyze an analogue of the Erdős-Füredi construction that satisfies an additional restriction on scalar products.

4 Acknowledgements

The author is very grateful to János Pach, Sasha Polyanskii and Konrad Swanepoel for several stimulating discussions on the subject and to one of the reviewers, who had very carefully read the manuscript and pointed out several problems with the proofs and the exposition. It helped to improve a lot the presentation of the paper.

References

- [1] L. Danzer, B. Grünbaum, *Über zwei Probleme bezüglich konvexer Körper von P. Erdős und von V. L. Klee*, Math. Zeitschrift 79 (1962), 95–99.
- [2] P. Erdős, *Some unsolved problems*, Michigan Math. J. 4 (1957), 291–300.
- [3] P. Erdős, Z. Füredi, *The greatest angle among n points in the d -dimensional Euclidean space*, Annals of Discrete Math. 17 (1983), 275–283.
- [4] V. Harangi, *Acute sets in Euclidean spaces*, SIAM J. Discrete Math. 25 (2011), N3, 1212–1229.
- [5] H. Martini, V. Soltan, *Antipodality properties of finite sets in Euclidean space*, Discrete Mathematics 290 (2005), 221–228.
- [6] J. Pach, *A remark on transversal numbers*, In: The mathematics of Paul Erdős II, eds. R. L. Graham et al., Algorithms and Combinatorics, 14, Springer, Berlin (1997), 310–317.
- [7] J. Pach, K. Swanepoel, *Double-normal pairs in space*, Mathematika 61 (2015), N1, 259–272. arXiv:1404.0419v1
- [8] J. Pach, K. Swanepoel, *Double-normal pairs in the plane and on the sphere*, Beitrage zur Algebra und Geometrie 56 (2015), N2, 423–438. arXiv:1404.2624v1